

Effects of Bacterial Strains to Inhibit Growth of *Phytophthora pistaciae* under Different Electrical Conductivities

Moslem Hajabdolahi¹, Mohammad Moradi^{*2,3}, Seyed Reza Fani⁴

¹Department of Plant Pathology, Faculty of Agriculture, Islamic Azad University, Damghan Branch, Damghan, Iran

²Pistachio Research Center, Horticultural Sciences Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Rafsanjan, Iran

³Pistachio Safety Research Center, Rafsanjan University of Medical Sciences, Rafsanjan, Iran

⁴Plant Protection Research Department, Yazd Agricultural and Natural Resources Research and Education Center, AREEO, Yazd, Iran

Received: 7 May 2017

Accepted: 16 April 2018

Abstract

Root and crown rot (gummosis) is known as the most destructive disease affecting pistachio in Iran. The efficiency of bacterial strains to reduce the growth rate of *Phytophthora pistaciae* was studied under different electrical conductivities (EC, 0, 2, 4, 8, 12 ds/m). Soil and rhizosphere samples were collected from pistachio growing regions in Kerman province, Iran, during 2011 - 2012. Overall, the strains of bacteria were presented in all sampling areas in both infected and uninfected orchards. Out of 400 bacterial isolates, 63% and 37% were collected from soil and rhizosphere samples, respectively. Among 400 bacterial isolates, 19 exhibited the highest ability to reduce the growth of *P. pistaciae* in dual culture, volatile and non-volatile compounds, though by different degrees. The degrees of inhibitory activities against mycelial growth of *P. pistaciae* by *Pseudomonas fluorescens* strains ranged from 40 to 97.5%, 8 to 97.5% and 7.5 to 90% in dual culture, non-volatile and volatile assays, respectively. The *Bacillus subtilis* strains reduced the growth of *P. pistaciae* by 22-92.5%, 17-85%, 21-92.5% in dual culture, non-volatile and volatile assays, respectively. The negative effects of ECs on the growth of *P. pistaciae* in modified CMA were observed in 8 and 12 ECs. ECs had no effect until 8 ds/m on the growth of *P. pistaciae*, while the mycelial growth decreased by ECs higher than 8 ds/m. No mycelial growth was observed at EC 14 ds/m. There were significant differences between different bacterial isolates, ECs and their interactions on the mycelial growth of *P. pistaciae*. The highest mycelial suppression belonged to isolates Nos. 123 and 112 in dual culture, volatile and non-volatile compounds test. More research is required to understand the native mechanisms involved in biological control under natural conditions in pistachio orchards.

Keywords: Biological control, Electrical conductivities, *Phytophthora*, Rhizosphere, Salinity, Sustainable agriculture.

Introduction

Both biotic and abiotic factors may affect pistachio (*Pistacia vera* L.) production (Moradi, 2015; Moradi *et al.*, 2017). Crown and root rot in pistachio

trees (gummosis) are caused by multiple species of the *Phytophthora* (Ershad, 1992; Banihashemi, 1994; Banihashemi and Moradi, 2004; Mirabolfathy *et al.*,

*Corresponding author: Email: moradi@pri.ir

2001; Fani *et al.*, 2005; Mostowfizadeh-Ghalamfarsa *et al.*, 2008). *Phytophthora pistaciae* and *P. drechsleri* are the most frequently found species (Mirabolfathy *et al.*, 2001; Banihashemi and Moradi, 2004). Pistachio gummosis is the most common cause of tree mortality in the orchards, leading to a remarkable crop reduction. Annual damage in terms of tree mortality is estimated to be between 2 to 11% (Moradi, 2015; Moradi *et al.*, 2017). Different management strategies have been introduced including the use of resistant rootstocks, the application of cultural practice, the use of chemical pesticides as well as biological approaches (Moradi, 2015; Moradi *et al.* 2017; Fani *et al.*, 2013). Mutualism of rhizobacteria or plant growth-promoting rhizobacteria (PGPR) and plants is a key factor in the biocontrol strategies for an ecofriendly management of plant diseases in many plant-microbe interactions. Disease is suppressed through induced systematic resistance (ISR) and through the production of antifungal metabolites (Vessy, 2003). *In vitro* studies have shown that different metabolites such as antibiotics, enzymes, and volatile compounds produced by *Bacillus* and *Pseudomonas* isolates are the main reasons for the phytopathogen suppression ((Haas and Défago, 2005; Lugtenberg and Kamilova, 2009). Antagonism and plant growth promotion effects of the metabolites on plant pathogens are demonstrated in natural environments (Earl *et al.*, 2008). Overall, plants exposed to environmental stresses are more susceptible to root rot caused by *Phytophthora* root rot. On the other hand, environmental stress increased the severity and incidence of plant disease in most cases. Swiecki and MacDonald (1991) have shown higher root and crown rot severity in stressed tomato plants caused by *Phytophthora parasitica* than non-stressed controls. The synergistic effects were the most pronounced in young (pre-bloom) plants. Besri (1993) reported that the isolates of *Phytophthora* gummosis of citrus were more tolerant to soil salinity than the crop plant they attacked. The isolates produce more sporangia in saline soils than in non-saline ones (Besri 1993). It has been

shown that increasing the salinity of media promotes mycelial growth of *P. citrophthora* and *P. parasitica* with an optimum of -1.44 to -3.11 bars (Benyahia, 1998) as well as the stimulation of sporangium formation of *Phytophthora parasitica* by salinity under *in vitro* conditions (Regragui and Lahlou, 2005). Banihashemi and Tabatabaee (2004) reported that increasing the salinity had significant effects on the growth of *P. citrophthora* and *P. nicotianae* while mycelial growth, sporangia and zoospores of *P. nicotianae* were more sensitive than *P. citrophthora*. There is not enough information on the interactions between profitable bacterial strains of pistachio rhizosphere and *Phytophthora* species in the salinity status. Therefore, the present study was conducted to evaluate the capability of bacterial isolates to inhibit *P. pistaciae* under ECs conditions.

Materials and Methods

Bacterial isolates

A total of 130 samples were collected either from soil or rhizosphere of pistachio producing areas in Kerman province during 2010-2012. Samples were taken according to a diagonal pattern in each orchard (n=24 subsamples). Samples were thoroughly homogenized in the laboratory to obtain three 100 g subsamples. Serial dilution method was used to isolate the bacteria from soil and rhizosphere on nutrient agar medium (NA, Himedia Pvt. Ltd., India). Petri-plates were incubated at 28°C for 48 hours. The suspected colonies (in terms of shape, color and size) were sub-cultured to a new NA plates to obtain single colonies to study antagonistic effects.

P. pistaciae isolate

Originally obtained from infected trees in Rafsanjan region deposited in the culture collection at the Iranian Research Institute of Plant Protection (IRIPP), a virulent strain of *P. pistaciae* was used in all experiments. The isolate was re-cultured and maintained on either Potato Dextrose Agar (PDA, Merck, Germany)

or Corn Meal Agar (CMA) (Himedia, Pvt. Ltd., India) before use.

Screening of bacterial antagonistic

Abilities of isolated bacteria to reduce mycelial growth of *P. pistaciae* were assessed in dual culture (Aghighi *et al.*, 2004), volatile compounds (Hora and Baker, 1972) and non-volatile compounds (Berg and Ballin, 1995) assays.

Effect of salinity on mycelial growth of *P. pistaciae*

The ability of selected bacterial strains to withstand two salt compounds (NaCl and CaCl₂) was tested in different electrical conductivities (ECs). Briefly, modified corn meal agar (MCMA) medium was prepared with different levels of NaCl and CaCl₂ salts in the range of 0, 2, 4, 8, 10, 12, 14 and 20 ds/m according to Richards (1954), and was measured using an electrical conductivity meter.

To evaluate the mycelium growth inhibition, a 5 mm agar plug of *P. pistaciae* was placed in the center of a petri dish plate containing the MCMA medium, and incubated at 27°C in a dark environment. The plates were monitored every 48 hours for 7 days to measure mycelial growth and colony diameters (Aghighi *et al.* 2004).

Effect of salinity on the antagonistic efficiency of bacterial strains

The effect of salinity on the antagonistic efficiency of bacterial strains to reduce mycelial growth of *P. pistaciae* was assessed through dual culture, volatile and non-volatile compounds in modified CMA medium, as already described (Aghighi *et al.*, 2004; Hora and Baker, 1972; Berg and Ballin, 1995).

The ability of the bacterial strains to inhibit fungal growth of *P. pistaciae* was calculated using the following formula:

$$I = (C - T) / C \times 100$$

where *I* is the inhibition percentage of mycelial growth, *C* is the mycelial growth in control plates and *T* is mycelial growth in the treatments.

Identification of bacterial strains

The nineteen selected bacterial strains with the highest antagonistic activity were identified based on morphological and biochemical characteristics (Shaad *et al.*, 2001).

Statistical analysis

The experiments were carried out based on completely randomized design with three replicates. The average values of mycelial growth were separately calculated for each replication. Mean comparisons were done using Duncan's new multiple range test at 1 % probability. If needed, data was log-transferred prior to the analysis.

Results

Overall, bacteria strains were isolated in all sampling areas although their frequencies were different in both infected and uninfected orchards. Out of 400 bacterial isolates, 63% and 37% were collected from soil and rhizosphere samples, respectively. Among 400 bacterial isolates, 170 were able to inhibit the growth of *P. pistaciae* in screening through dual culture assays. Out of 170 isolates, 19 had shown the highest ability to reduce growth of *P. pistaciae* in dual culture, volatile and non-volatile compounds (Table 1). Bacterial strains with the highest antagonistic potential were identified based on biochemical and morphological characterizations (Table 2).

The intensity of inhibitory activities against mycelial growth of *P. pistaciae* by *P. fluorescens* ranged from 40 to 97.5%, 8 to 97.5% and 7.5 to 90% in dual culture, non-volatile and volatile assays, respectively (Table 1). The *Bacillus subtilis* strains reduced the growth of *P. pistaciae* by 22-92.5, 17-85, 21-92.5 in dual culture, non-volatile and volatile assays, respectively. Among the 19 isolates, isolates Nos. 123 and 112 showed the highest ability to reduce mycelial

growth of *P. pistaciae* compared to other isolates. As shown in Table 2, the growth of *P. pistaciae* was not significantly affected by 2 and 4ECs (ds/m) compared to the control in CMA medium either in presence or absence of bacterial isolates. On the other hand, the effects of ECs in modified CMA were significant in 8 and 12ECs (ds/m), which reduced the mycelial growth of *P. pistaciae* ranged respectively from 38 to 92.5% and 40 to 97.5% in dual culture, 31.5 to 92 and 37.5 to 97.5 in non-volatile compounds and 34 to 83.5 and

7.5 to 92.5 in volatile compounds assays. The growths of *P. pistaciae* in different ECs have shown no effects until 8 ds/m while the mycelial growth decreased in the ECs higher than 8 ds/m. No mycelial growth was observed at ECs 14 ds/m. There were significant differences between the isolates of bacteria, ECs and their interactions to reduce mycelial growth of *P. pistaciae* (Table 3). The highest mycelial reductions belonged to isolates Nos. 123 and 112 in dual culture, volatile and non-volatile compound tests.

Table1. The effect of bacterial strains on mycelial growth (CM) of *Phytophthora pistaciae* in different electrical conductivities by various competition tests

	Ec	Bacterial strains																			
		123	32	28	112	146	151	23	153	154	143	150	29	79	149	6	78	157	40	156	Control
Dual Culture	0	0.9	1.9	2.3	0.9	3.9	4.0	3.4	3.7	1.3	1.3	1.6	1.3	4.5	4.3	6.7	1.7	5.0	1.9	4.1	8.6
	2	0.7	1.5	2.0	0.8	3.6	4.0	3.3	3.6	1.3	1.3	1.5	1.3	4.0	3.8	4.7	1.6	3.0	1.7	3.8	8.5
	4	0.7	1.7	2.0	0.8	3.4	3.6	3.3	3.5	1.2	1.2	1.3	1.2	3.7	3.6	4.4	1.3	3.3	1.5	3.7	8.7
	8	0.4	1.2	1.4	0.4	2.3	2.4	1.9	1.7	0.8	0.8	1.0	0.8	2.9	2.5	2.9	1.2	4.5	1.4	2.4	7.3
	12	0.1	0.5	2.4	0.1	1.3	1.4	1.0	1.0	0.3	0.3	0.5	0.3	1.5	1.3	1.6	0.5	2.2	0.7	1.4	4.0
	LSD	0.2	0.5	2.6	0.2	0.5	0.5	0.3	0.4	0.4	0.3	0.2	0.3	0.3	0.6	0.6	0.4	0.2	0.8	0.4	0.6
Non-Volatile compounds	0	1.3	2.6	3.8	1.3	5.8	5.4	4.8	5.2	2.8	2.4	2.7	7.9	4.5	5.5	7.1	2.5	5.5	2.8	4.9	8.6
	2	1.5	2.6	2.8	1.8	4.0	5.2	4.5	5.2	2.8	2.4	2.5	7.3	4.0	5.1	5.7	2.4	3.7	2.5	4.0	8.5
	4	1.8	2.7	3.5	1.6	4.1	4.9	4.3	4.9	2.7	2.4	2.4	7.0	3.7	4.8	5.4	2.4	3.7	2.4	4.0	8.7
	8	0.6	1.7	2.1	0.8	2.8	2.9	2.7	2.4	1.8	1.7	1.7	5.0	2.9	3.2	3.5	1.6	4.8	1.5	2.7	7.3
	12	0.1	0.9	1.1	0.2	1.5	1.4	1.2	1.2	0.7	0.5	0.6	2.5	1.5	1.8	1.8	0.7	2.4	0.9	1.5	4.0
	LSD	0.5	0.3	0.6	0.2	0.5	0.3	0.5	0.5	0.5	0.2	0.3	0.4	0.6	0.8	0.5	0.3	0.4	0.3	0.3	0.3
Volatile compounds	0	1.2	2.3	3.7	2.6	5.2	5.0	4.7	4.6	2.6	2.3	2.7	7.7	4.5	5.3	6.8	2.2	5.1	2.4	4.5	8.6
	2	1.2	2.3	3.7	2.6	3.5	4.9	4.0	5.0	2.5	2.2	2.3	7.1	3.8	4.7	5.4	2.1	3.2	2.4	3.8	8.5
	4	1.2	2.3	3.7	2.6	3.4	4.7	4.1	4.7	2.5	2.2	2.2	6.8	3.6	4.3	5.2	2.2	3.4	2.2	3.9	8.7
	8	1.2	2.3	3.7	2.6	2.5	2.8	2.2	2.1	1.5	1.5	1.6	4.8	2.5	2.8	3.1	1.4	4.6	1.4	2.5	7.3
	12	1.2	2.3	3.7	2.6	1.3	1.2	1.0	1.0	0.3	0.4	0.5	2.3	1.3	1.4	1.5	0.6	2.1	0.7	1.4	4.0
	LSD	0	0	0	0	0.4	0.2	0.4	0.5	0.5	0.3	0.3	0.4	0.5	0.8	0.5	0.3	0.3	0.5	0.3	0.3

Table 2. Characterizations of bacterial strains and their efficiency on *P. pistaciae* mycelial growth reduction

No.	IPRC code	Bacterial species	Location	Sampling area	Orchard contamination to gummosis	Mycelial growth reduction rate (%)		
						Dual culture	Non-Volatile compounds	Volatile compounds
1	23	<i>Pseudomonas fluorescens</i>	Tajabad (Rafsanjan)	Rhizosphere	-	60-75	44-70	45-75
2	28	<i>Pseudomonas fluorescens</i>	Esmailabad (Nough)	Rhizosphere	-	40-81	56-72.5	7.5-57
3	29	<i>Pseudomonas fluorescens</i>	Esmailabad (Nough)	Soil	+	85-92.5	8-37.5	10-42.5
4	32	<i>Pseudomonas fluorescens</i>	Tajabad (Rafsanjan)	Rhizosphere	+	69-77.5	69-77.5	42.5-73.5
5	79	<i>Pseudomonas fluorescens</i>	Khalilabad	Rhizosphere	+	48-62.5	48-62.5	48-67.5
6	112	<i>Pseudomonas fluorescens</i>	Sirjan	Rhizosphere	+	89.5-97.5	79-95	35-70
7	123	<i>Pseudomonas fluorescens</i>	Sirjan	Rhizosphere	-	89.5-97.5	79-97.5	70-86
8	143	<i>Pseudomonas fluorescens</i>	Tajabad (Rafsanjan)	Rhizosphere	+	85-92.5	72-87.5	73-90
9	146	<i>Pseudomonas fluorescens</i>	Kerman	Soil	-	55-68	32.5-62.5	39.5-67.5
10	149	<i>Pseudomonas fluorescens</i>	Kerman	Soil	-	50-67.5	36-56	38-65
11	151	<i>Pseudomonas fluorescens</i>	Kerman	Rhizosphere	+	53-67	37-65	42-70
12	153	<i>Pseudomonas fluorescens</i>	Kerman	Rhizosphere	-	57-77	39-70	41-75
13	6	<i>Bacillus subtilis</i>	Kousarriz (Rafsanjan)	Rhizospher	+	22-60	17-55	21-62.5
14	40	<i>Bacillus subtilis</i>	Anar	Rhizosphere	-	78-83	67-79	72-82.5
15	78	<i>Bacillus subtilis</i>	Khalilabad	Rhizosphere	-	80-87.5	71-82.5	74-85
16	150	<i>Bacillus subtilis</i>	Kerman	Soil	-	81-87.5	69-85	69-87.5
17	154	<i>Bacillus subtilis</i>	Kerman	Soil	+	85-92.5	67-82.5	70-92.5
18	156	<i>Bacillus subtilis</i>	Kerman	Soil	-	52-67	43-63	48-66
19	157	<i>Bacillus subtilis</i>	Kerman	Soil	-	38-65	34-57	37-62

Table 3. Analysis of variance of growth inhibition of *P. pistaciae* under *in vitro* experiments.

Source	Treatment	df	sum of square	Mean of Square	P
Dual culture	Bacteria	17	739.47	38.91	<.0001
	EC	4	156.87	39.21	<.0001
	Bacteria * EC	76	85.88	1.13	<.0001
Non-volatile	Bacteria	17	693.53	36.5	<.0001
	EC	4	362.56	90.64	<.0001
	Bacteria * EC	76	93.86	1.23	<.0001
Volatile	Bacteria	17	622.85	32.78	<.0001
	EC	4	271.03	67.75	<.0001
	Bacteria * EC	76	138.93	1.828	<.0001

Discussion

One of the major factors reducing pistachio quality and quantity is the increase in the salinity of soil and water in recent years. Electrical conductivity in soil and water is about 2.0 to more than 20 dS/m, which have been showing an increasing trend in recent years because of dropping underground water table and as a consequence of water quality decrease. The current study has shown the presence of native strains of bacterial with an ability to control *Phytophthora* crown and root rot of pistachio trees under saline and non-saline conditions. The bacterial isolates are already adapted to biotic and abiotic stresses found in the orchards and may have a higher stability to control pistachio gummosis. This requires more research to understand the dynamics of bacterial isolates in soil and rhizosphere (Lee *et al.*, 2015).

The effects of different ECs levels were compared on the mycelial growth of *P. pistaciae* in a modified medium. No effect was observed on the mycelial growth of *P. pistaciae* in 4 and 8 ECs while a salinity of higher than 8 ds/m decreased the mycelial growth to different degrees. Here, it is shown that the bacterial isolates could inhibit mycelia growth of *P. pistaciae* either in the presence or absence of salt stresses, which was more pronounced in higher concentrations of salt in the media. Several studies have shown an increase of mycelial growth and spore germination by chloride salts in low concentrations, while high concentrations inhibit the growth and spore germination of many plant pathogenic fungi compared to the con-

trol – although by different degrees (Benyahia, 1998; Biggs *et al.*, 1997; Regragui and Lahlou, 2005). The effects of salts in reducing growth of fungal species may be due to toxicity in high concentrations affecting the osmotic balance or pressure in fungal cells, which inhibit biological activity (Arras *et al.*, 1998). On the other hand, in low concentrations, the effect of salinity to stimulate the growth of fungal species is under the effect of specific ions (Wisniewski *et al.*, 1995). In the present study, there were significant differences between isolated bacteria to reduce the growth of *P. pistachio* with the highest effects, which were observed in *Pseudomonas fluorescens* isolates Nos. 112 and 123 in all *in vitro* assays. Raaijmakers *et al.*, (2010) have shown that the production of volatile and non-volatile compounds is associated with the inhibition of pathogens in the rhizosphere of plants. The production of metabolites such as antibiotics and enzymes in *Pseudomonas fluorescens* and *Bacillus* isolates with a wide range of activities may act as antagonistic factors in reducing the growth of *P. pistaciae*. This has already been shown by other researchers (Haas and Défago, 2005; Lugtenberg and Kamilova, 2009; Raaijmakers *et al.*, 2002; Raaijmakers and Mazzola, 2012; Shaad *et al.*, 2001). Understanding the population dynamics and diversity of bacteria associated with rhizosphere of pistachio may be useful to improve plant growth due to the synthesis of phytohormones, such as IAA, siderophores, ACC deaminase, solubilizing of Phosphate, and improve-

ment of mineral uptake (Lee *et al.*, 2015; Compant *et al.*, 2010; Kloepper *et al.*, 1989; Asghar *et al.*, 2004; Glick *et al.*, 2007) as well as biotic and abiotic stresses, in particular. Studies have been carried out on the effects of salinity on the growth of different fungal pathogens. For example, Boumaaza *et al.* (2015) showed that the salinity (NaCl and CaCl₂ concentrations) stimulated the growth of *Botrytis cinerea* isolates up to 150ppm and higher concentrations reduced the mycelial growth and inhibit spore germination while increasing the formation of sporangia in all concentrations. Benyahya (1998) demonstrated that increasing the salinity promotes *in vitro* mycelial growth of *P. citrophthora* and *P. parasitica*, the causal agents of root rot of citrus. In view of our findings, no effects was observed at low concentrations on the mycelial growth of *P. pistaciae* (4 and 8 dS/m), while higher salinity (higher than 8 dS/m) reduced the growth to different degrees compared to the control. More research is required to understand the native mechanisms involved in biological control, the effects of specific ions and their concentrations on the growth and sporulation of *Phytophthora* spp. as well as the severity of crown and root rot of pistachio trees under natural conditions in pistachio orchards. Recent observations have shown that the applications of calcium sulfate in pistachio orchards could suppress the incidence and severity of crown and root rot of pistachio trees. Several studies have showed that applications of calcium salts can suppress diseases caused by several pathogens (Biggs *et al.*, 1997; Conway *et al.*, 1992; Volpin and Elad, 1991; Yamazaki and Hoshina, 1995).

Acknowledgements

The authors are grateful to the staff of Pistachio Research Center involved, to A. Farajpour for his cooperation and assistance.

References

- Aghighi S, Shahidi Bonjar GH, Rawashdeh R, Batayneh S, Saadoun I (2004) First report of antifungal spectra of activity of Iranian *Actinomyces* strains against *Alternaria solani*, *Alternaria alternata*, *Fusarium solani*, *Phytophthora megasperma*, *Verticillium dahliae* and *Saccharomyces cerevisiae*. Asian Journal of Plant Science. 3(4), 463-471.
- Arras G, Sanna P, Astone V, Arru S (1998) Effect of calcium chloride on the inhibitory activity of *Rhodotorula glutinis* against *Penicillium italicum* on orange fruits. Italus Hortus. 5, 67-70.
- Asghar HN, Zahir ZA, Arshad M (2004) Screening rhizobacteria for improving the growth, yield, and oil content of canola (*Brassica napus* L.). Crop and Pasture Science. 55(2), 187-194.
- Banihashemi Z, Moradi M (2004) The Frequency of isolation of *Phytophthora* spp. from crown and root of pistachio nut tree and reaction of the crown and root to the casual agents. Iranian Journal of Plant Pathology. 40, 57-77.
- Banihashemi Z, Tabatabaee SAR (2004) Interaction between salinity and *Phytophthora citrophthora* in pistachio seedlings under hydroponic system. Iranian Journal of Plant Pathology. 40, 159-178.
- Berg G, Ballin G (1994) Bacterial antagonists to *Verticillium dahliae* Kleb. Journal of Phytopathology. 141(1), 99-110.
- Besri M (1993) Effects of salinity on plant diseases development. In: Lieth H, Al Masoom A (eds) Towards the rational use of high salinity tolerant plants. Springer Science & Business Media.
- Biggs AR, El-Kholi MM, El-Neshawy S, Nickerson R (1997) Effects of calcium salts on growth, polygalacturonase activity, and infection of peach fruit by *Monilinia fructicola*. Plant Disease. 81(4), 399-403.

- Boumaaza B, Benkhelifa M, Belkhouja M (2015) Effects of two salts compounds on mycelial growth, sporulation, and spore germination of six isolates of *Botrytis cinerea* in the western north of Algeria. International Journal of Microbiology. <https://www.hindawi.com/journals>. doi: 10.1155/2015/572626.
- Compant S, Clément C, Sessitsch A (2010) Plant growth-promoting bacteria in the rhizo- and endosphere of plants: their role, colonization, mechanisms involved and prospects for utilization. Soil Biology and Biochemistry. 42(5), 669-678.
- Conway WS, Sams CE, McGuire RG, Kelman A (1992) Calcium treatment of apples and potatoes to reduce postharvest decay. Plant Disease. 76(4), 329-334.
- Fani SR, Moradi M, Alipour MM, Sherafati A, Mohammadi MM, Sedaghati E, Khodaygan P (2013) Efficacy of native strains of *Trichoderma harzianum* in biocontrol of pistachio gummosis. Iranian Journal Plant Protection Science. 44(2), 243- 252. [In Persian].
- Fani SR, Zamanizadeh HR, Mirabolfathy M (2005) Isolation and identification of the causal agents of root and crown rot of pistachio trees in the Sistan and Baluchistan provinces. Acta Horticulture. 726, 647-650.
- Glick BR, Cheng Z, Czarny J, Duan J (2007) Promotion of plant growth by ACC deaminase-producing soil bacteria. European Journal of Plant Pathology. 119(3), 329-339.
- Haas D, Défago G (2005) Biological control of soil-borne pathogens by fluorescent pseudomonads. Nature Reviews Microbiology. 3(4), 307-319.
- Hora TS, Baker R (1972) Soil fungistasis: microflora producing a volatile inhibitor. Transactions of the British Mycological Society. 59(3), 491-500.
- Kloepper JW, Lifshitz R, Zablutowicz RM (1989) Free-living bacterial inocula for enhancing crop productivity. Trends Biotechnology. 7(2), 39-44.
- Lee BD, Dutta S, Ryu H, Yoo SJ, Suh DS, Park K (2015) Induction of systemic resistance in *Panax ginseng* against *Phytophthora cactorum* by native *Bacillus amyloliquefaciens* HK34. Journal of Ginseng Research. 39(3), 213-220.
- Lugtenberg B, Kamilova F (2009) Plant-growth-promoting rhizobacteria. Annual Review of Microbiology 63, 541-556.
- Mirabolfathy M, Cooke D, Duncan JM, Williams NA, Ershad D, Alizadeh A (2001) *Phytophthora pistaciae* sp. nov. and *P. melonis*: the principal causes of pistachio gummosis in Iran. Mycological Research. 105, 1166-1175.
- Moradi M (2015) Assessment of application of systemic and protective fungicides for long-term control of pistachio crown and root rot. Final Report of Iranian Pistachio Research Institute 2-06-06-88008. ACIST Register number: 47569. [In Persian].
- Moradi M, Mohammadi AH, Haghdel M (2017) Efficiency of Elite Fungicide for Control of Pistachio Gummosis. Journal of Nuts. 8(1), 11-20.
- Raaijmakers JM, De Bruijn I, Nybroe O, Ongena M (2010) Natural functions of lipopeptides from *Bacillus* and *Pseudomonas*: more than surfactants and antibiotics. FEMS Microbiology Reviews. 34(6), 1037-1062.
- Raaijmakers JM, Mazzola M (2012) Diversity and natural functions of antibiotics produced by beneficial and plant pathogenic bacteria. Annual Review of Phytopathology. 50, 403-424.
- Raaijmakers JM, Vlami M, De Souza JT (2002) Antibiotic production by bacterial biocontrol

- agents. *Antonie van Leeuwenhoek*. 81(1-4), 537-547.
- Regragui A, Lahlou H (2005) Effect of salinity on in vitro *Trichoderma harzianum* antagonism against *Verticillium dahliae*. *Pakistan Journal of Biological Sciences* 8(6), 872-876.
- Richards LA (1954) Diagnosis and improvement of saline and alkali soils. Washington: United States Salinity Laboratory, United States Department Of Agriculture; Washington. pp 169.
- Shaad NW, Jones JB, Chun W (2001) Laboratory Guide for Identification of Plant Pathogenic Bacteria, 3rd ed., American Phytopathological Society, APS Press, St. Paul, MN, USA, pp 398.
- Swiecki TJ, MacDonald JD (1991) Soil salinity enhances *Phytophthora* root rot of tomato but hinders asexual reproduction by *Phytophthora parasitica*. *Journal of the American Society for Horticultural Science*. 116(3), 471-477.
- Vessy K (2003) Plant Growth Promoting Rhizobacteria as Biofertilizers. *Plant and Soil*. 255(2), 571–586.
- Volpin H, Elad Y (1991) Influence of calcium nutrition on susceptibility of rose flowers to *Botrytis* blight. *Phytopathology*. 81(11), 1390-1394.
- Wisniewski M, Droby S, Chalutz E, Eilam Y (1995) Effects of Ca^{2+} and Mg^{2+} on *Botrytis cinerea* and *Penicillium expansum* in vitro and on the biocontrol activity of *Candida oleophila*, *Plant Pathology*. 44, 1016–1024.
- Yamazaki H, Hoshina T (1995) Calcium nutrition affects resistance of tomato seedlings to bacterial wilt. *HortScience*. 30(1), 91-93.